

# Chemistry 110 Lecture Unit 5

## Chapter 11-GASES

### I. PROPERTIES OF GASES

- A. Gases have an indefinite shape.
- B. Gases have a low density
- C. Gases are very compressible
- D. Gases exert pressure equally in all directions on the walls of a container.
- E. Gases mix spontaneously and completely with one or more other gases.

### II. KINETIC MOLECULAR THEORY

- A. Gases are composed of such extremely tiny atoms or molecules that are widely separated by empty space.
- B. Gas particles move in a random, rapid, and continuous motion, thus has kinetic energy.
- C. Gas particles move so rapidly and are so far apart the there is essentially no force of attraction between the particles.
- D. Particles collide frequently with each other and with the walls of the container, the collisions are perfectly "elastic" - (No net loss of energy as a result of a collision)

### III. AVERAGE KINETIC ENERGY

The average kinetic energy (energy of motion) of the gas particles are directly proportional to its absolute  $T^{\circ}$  (Kelvin)

### IV GRAHAM'S LAW OF EFFUSION

The rate of diffusion of a gas is inversely proportional to it's size [Molar Mass]

### V. GAS MEASUREMENTS

#### A. Pressure

1. Pressure =  $\frac{\text{force}}{\text{Unit area}}$

2. Gases exert pressure equally in all directions on wall of a container.

3. Units

a) Types

Pascal	mm Hg	torr	Atmosphere	Psi
Pa			atm	

b) Conversions

*KNOW*  $\rightarrow 1 \text{ atm} = 760 \text{ mm Hg} = 760 \text{ torr}$  (exactly)

$1.013 \times 10^5 \text{ Pa} = 1 \text{ atm} = 14.68 \text{ psi}$

## B. Temperature

Absolute temperature (Kelvin) Conversions

$$K = ^\circ C + 273.15$$

## C. Volume

1. The volume of a gas is the volume of the container it occupies.
2. Units: liters or milliliters

## V. RELATIONSHIP BETWEEN $^{\circ}T$ , VOLUME, AND PRESSURE.

### A. Boyle's law P & V

As the pressure increases the volume decreases in the same proportion.

### B. Charles's law $^{\circ}T$ & V

As the temperature (Kelvin) is increased the volume is increased proportionally.

C. Gay-Lussac's Law

When temperature (K) increases pressure increases proportionally.

D. COMBINED GAS LAW

P, V, and °T varying. Assume that the mass is constant.

Problems:

1.

$$P_1 = 3.0 \text{ atm}$$

$$T_1 = 2 \text{ }^\circ\text{C}$$

$$V_1 = 29 \text{ L}$$

$$P_2 = ?$$

$$T_2 = 300.0 \text{ K}$$

$$V_2 = 100.0 \text{ L}$$

2. A certain mass of gas occupies 5.50 L at 34°C and 655 mm Hg. What will its volume in liters be if it is cooled to 10.0°C and its pressure doubles.

### E. GAY-LUSSAC'S LAW OF COMBINING VOLUMES

At the same °T and Pressure, the volumes of gases that combine in a chemical reaction are in the ratio of small whole numbers.

### F. MOLAR GAS VOLUME; AVOGARDO'S HYPOTHESIS

At the same temperature and pressure the same number of moles of different gases have the same volume. The *Molar Volume* is the volume of one mole of any gas at a given °T & P. [STP]

Standard temperature and Pressure = [STP]:

1 mole of gas = 22.4 L At: 273 K and  
1 atm (760 torr)

Conversion factors:  $\frac{1 \text{ mole}}{22.4 \text{ L}}$  ;  $\frac{22.4 \text{ L}}{1 \text{ mole}}$

\*Warning: Use  
only with STP!

#### Calculations using STP

1. A 2.00 L sample of a gas at 0°C and 1.00 atm has a mass of 3.94g. Calculate.....

a) Density

b) Molar mass

2. What is the density of ammonia gas at 273K and 760 torr?

3. The density of an unknown gas is 1.43 g/L at 0 ° C and 760 torr. What is the molar mass of the unknown gas?

G. IDEAL GAS EQUATION:

Derivation:

KNOW:  $PV=nRT$       Where: n = moles of gas

$$R = \frac{0.0821 \text{ L-atm}}{\text{mole-K}}$$

H. Problems using  $PV=nRT$

1. What volume in liters will be occupied by 6.00 mol carbon dioxide gas at 105 mm Hg and 28°C?

2. What is the temperature in degrees Celsius of 0.500 mole hydrogen gas in a 463 mL container at a pressure of 0.500 atm?

### I. Molar Mass Calculations using $PV=nRT$

An 2.00 L sample at an unknown gas had a pressure of 755 mm Hg and temperature of 259 K. The mass of the sample was 3.94 g. What is it's molar mass?

#### WHEN TO USE:

1)  $PV = nRT$

mole

2) 22.4 L at STP

3) 
$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

#### Problems:

1. Calculate the molar mass of nitrogen gas at 0 °C and 1 atm, if the density is 1.25 g/L.

2. How many grams of ammonia gas in a 3999 ml sample at 9.8 atm and 20.0 °C?



3. At  $45^{\circ}\text{C}$  and  $1.20\text{ atm}$  the volume of  $1390\text{ mg}$  of fluorine gas is  $794\text{ mls}$ . Use this data to calculate the molar mass of fluorine gas.

4. A balloon has a volume of  $2.50\text{ L}$  at  $25^{\circ}\text{C}$ . When the balloon is placed in a refrigerator, its volume decreases to  $2.33\text{ L}$  assuming the pressure is the same inside and outside the refrigerator, what is the temperature of the gas in the balloon inside the refrigerator in degrees Celsius?

5. A  $415\text{ ml}$  sample of gas in a steel cylinder has a pressure of  $3.29\text{ atm}$  and a temperature of  $125^{\circ}\text{C}$ . If the closed cylinder is cooled to  $20.0^{\circ}\text{C}$  at constant volume, what is the new pressure of the gas in atmospheres?

### **J. DALTON'S LAW OF PARTIAL PRESSURES; Mixtures of gases**

The total pressure of a mixture of gases is equal to the sum of the partial pressures exerted by each gas.

$$P_{\text{total}} = P_1 + P_2 + P_3 + \dots$$

Example: The total pressure in a 1.00 liter container is 725 mm Hg. The container contains water vapor and nitrogen gas.

If the partial pressure of the water vapor is 225 mm Hg, what is the partial pressure of the nitrogen gas.

$$P_{\text{total}} = P_{\text{N}_2} + P_{\text{H}_2\text{O}}$$

Problem: The partial pressures of a mixture of nitrogen, oxygen and carbon dioxide gases are, respectively, 325 mm Hg, 0.128 atm, and 159 mm Hg. What is the total pressure of the mixture of gases.

## K. GASES IN CHEMICAL REACTIONS (GAS STOICHIOMETRY)


Certain chemical reactions involve gas as a reactant or product. For these types of reactions, the stoichiometric calculations involve the use of:

1)  $PV = nRT$

2)  $22.4 \frac{\text{L}}{\text{mole}}$  at STP

3) Molar volumes

The general stoichiometric scheme



Problems:

1. How many liters of ammonia gas can be produced by the reaction of 735 ml hydrogen gas with an excess nitrogen gas at 425 °C and 135 atm?      nitrogen + hydrogen → ammonia

2. How many liters of carbon dioxide gas at 0 °C and 1 atm are produced by the complete combustion of 60.0 mol of liquid glucose,  $C_6H_{12}O_6$ ?

3. How many liters of the air pollutant  $NO(g)$  could be produced at 985 °C and a pressure of 30.0 atm by the reaction of oxygen gas with 455 g of nitrogen gas.

## ENERGY and HEAT CAPACITY CALCULATIONS CHAP 3.12

CHANGES IN TEMPERATURE AS A SUBSTANCE IS HEATED [Energy Added]

As a substance absorbs heat, the temperature rises. Different substances can absorb and store more heat than others.

ex. Al vs. water

heat vs. temperature

degrees vs. calories

Heat = mass x specific heat x change in temperature

$$Q = MC \Delta T$$

A. HEAT CAPACITY [Specific heat] - (Chapter 3 sec. 3.11)

The amount of heat required to raise the temperature of 1 g of a substance exactly 1°C.

How many calories are required to raise the temperature of 25 grams of water from 11 °C to 45 °C?

Example: How many degrees Celsius will the temperature rise if 25 g ether absorbs 160. cal of energy.

$$\text{Specific heat}_{\text{ether}} = \frac{0.529 \text{ cal}}{\text{g } ^\circ\text{C}}$$

## B. ENERGY AND CHANGE OF STATE (Back to chapter 12!)

Energy (as heat) is either lost or absorbed when a substance changes its state

Solid → Liquid

Liquid → Gas

Gas → liquid

C. HEAT OF VAPORIZATION- The quantity of heat needed to convert a liquid at its boiling point to the gaseous state.

Prob: How much heat is needed to convert 155 g water to steam at it's B.P.?

$$\Delta H_{\text{vap}} = \frac{2.26 \text{ KJ}}{\text{g}}$$

D. HEAT OF FUSION- The quantity of heat needed to convert a solid at its melting point to the liquid state.

Prob: How much energy is needed to convert 35 g of ice to water at it's M.P.?  $\Delta H_{\text{fus}} =$

$$\frac{3.35 \text{ J}}{\text{g}}$$

## E. CHANGES IN TEMPERATURE AND PHYSICAL STATE

Summary of  $T^{\circ}$  & State changes when Energy [heat] is added:

$$\text{Specific Heat} = \frac{\text{J}}{\text{g } ^{\circ}\text{C}} \text{ or } \frac{\text{cal}}{\text{g } ^{\circ}\text{C}} \text{ ...etc.}$$

$$\text{Heat of Fusion} = \frac{\text{KJ}}{\text{g}} \text{ or } \frac{\text{KJ}}{\text{mole}} \text{ or } \frac{\text{Kcal}}{\text{g}} \text{ ...etc.}$$

$$\text{Heat of Vaporization} = \frac{\text{KJ}}{\text{g}} \text{ or } \frac{\text{KJ}}{\text{mole}} \text{ or } \frac{\text{Kcal}}{\text{g}} \text{ ...etc.}$$

**Problem.** How much energy [Heat in kilojoules] is needed to convert 500.0 g of ice at  $-15.0^{\circ}\text{C}$  to steam at  $105.0^{\circ}\text{C}$ ?


$$H_{\text{fusion}} = 335 \text{ J/g} \quad \text{or} \quad 80 \text{ cal/g} \quad H_{\text{vap}} = 2.26 \text{ kJ/g} \quad \text{Or} \quad 540 \text{ cal/g}$$

$$\text{Specific heat of ice} = 2.10 \text{ J/g } ^{\circ}\text{C} \quad \text{or} \quad 0.50 \text{ cal/g}$$

$$\text{Specific heat of water} = 4.18 \text{ J/g } ^{\circ}\text{C} \quad \text{or} \quad 1.00 \text{ cal/g } ^{\circ}\text{C}$$

$$\text{Specific heat of steam} = 2.0 \text{ J/g } ^{\circ}\text{C} \quad \text{or} \quad 0.80 \text{ cal/g } ^{\circ}\text{C}$$

GRAPH:



Calculations:

**A → B**    HEATING A SOLID

**B → C**    SOLID → LIQUID

**C → D**    HEATING A LIQUID

**D → E**    LIQUID → GAS

**E → F**    HEATING A GAS

TOTAL HEAT ADDED



Problems:

1. Calculate the number of joules required to convert 58.9 g of ice at  $0^{\circ}\text{C}$  to water at  $81^{\circ}\text{C}$

2. Given a sample of 30.0 g of water at  $37^{\circ}\text{C}$ , calculate the quantity of heat in kilojoules that would be required to convert it to steam at  $100^{\circ}\text{C}$ .


# Chapter 12-LIQUIDS, SOLIDS, AND INTERMOLECULAR FORCES

## I. Interactions between Molecules

<b>Solid</b>	<b>Liquid</b>	<b>gas</b>
Strong attraction between species	Some attraction	No attraction between species
Shape		
Density		
Ability to flow		
Compressibility		

### A. EVAPORATION OF LIQUIDS

Evaporation is the change from the liquid state to the gaseous or vapor state at the surface.



### B. VAPOR PRESSURE OF LIQUIDS

The vapor pressure of a liquid is the pressure exerted by the vapor above a liquid.

Dynamic equilibrium : rate of evaporation = rate of condensation

Vapor Pressure of Water at various Temperatures	
Temperature (°C)	Vapor pressure (mm Hg)
0	4.63
10	9.2
20	17.5
30	31.8
40	55.3
50	92.5
60	149.4
70	233.7
80	355.1
90	525.8
100	760.0

### C. BOILING POINT

The boiling point of a liquid is the temperature at which a liquid is changed to a gas within the liquid (bubbles formed underneath the surface)

Variation of the Boiling Point of Water with Elevation		
Location	Elevation (feet above sea level)	Boiling Point (°C)
San Francisco, CA	0	100.0
Salt Lake City, UT	4,390	95.6
Denver, CO	5,280	95.0
La Paz, Bolivia	12,795	91.4
Mount Everest	28,028	76.5

### D. INTERMOLECULAR FORCES - Review!

The attractive forces between molecules

#### I. Types

a. Dipole-Dipole interaction:

Dipole - dipole interactions are electrostatic attractions between polar molecules

b. Hydrogen bonds:

A hydrogen bond is a relatively strong dipole-dipole attractive force between a hydrogen atom and a pair of nonbonding electrons on a F, O, or N atom

c. London forces

London forces are very weak electrostatic forces of attraction between molecules with "temporary" dipoles.

E. EFFECTS OF INTERMOLECULAR FORCES ON PROPERTIES OF LIQUIDS  
STRENGTHS OF INTERPARTICLE (MOLECULAR FORCES)

-----  
VAPOR PRESSURE

-----  
BOILING POINT

-----  
EVAPORATION

-----  
EXAMPLES:

1. Which has the higher boiling point....N<sub>2</sub> or H<sub>2</sub>S?


2. Which has the lowest vapor pressure...Water or PH<sub>3</sub>?

## PRACTICE EXAM 5

Show all work and significant figures must be correct


### PROBLEMS

1. For the following questions, identify the element whose atoms fit the following descriptions (**Use the periodic table**)
  - a. \_\_\_\_\_ Which has d electrons?  
a) hydrogen    b) Copper    c) nitrogen
  - b. \_\_\_\_\_ A metalloid in period 5
  - c. \_\_\_\_\_ The element in period 4, group IIIA
  - d. \_\_\_\_\_ The element with a total of 3 electrons in the 2<sup>nd</sup> main energy shell
  - e. \_\_\_\_\_ The element with 5 completely filled orbitals
  - f. \_\_\_\_\_ The smallest alkali metal
  - g. \_\_\_\_\_ The largest period 4 transition metal
  
- 2 a. Write the electron configuration and orbital diagram of a potassium ion
  
- b. Write the electron configuration and orbital diagram of tin
  
- 2) Which is larger?
  - a) B or F?
  - b) Mg or Sr?
  
- 3) Which Has more metallic properties?
  - a) Na or Cs?
  - b) Mn or Zn?
  
- 4) Is the following bonds (pure) covalent, polar covalent or ionic?
  - a) SO
  - b) HBr
  - c) KBr
  - d) Cl<sub>2</sub>
  
- 5) Are the following molecules (pure) covalent, polar covalent or ionic?
  - a) CO
  - b) O<sub>2</sub>
  - c) NO
  
7. Write the Lewis dot structure for the following compounds



8. The partial pressures of a mixture of nitrogen, oxygen, and carbon dioxide gases are, respectively, 325 mm Hg, 0.128 atm, and 159 mm Hg. What is the total pressure of this mixture of gases in torr?
9. The gas in an inflated balloon has a volume of 125 ml at 29 °C. What volume in liters will the gas in the balloon occupy at 137 °C if its pressure is held constant
10. A gas occupies a volume of 410.0 mL at 27°C and 740.0 mm Hg pressure. Calculate the volume the gas would occupy at STP.

11. Smelling salts contain ammonium carbonate, which can decompose to form ammonia, a mild heart stimulant. The ammonium carbonate decomposes according to the following reaction:




How many liters of  $\text{NH}_3$  at 25°C and 1.00 atm are formed from 0.500 g of  $\text{(NH}_4\text{)}_2\text{CO}_3$

12. A 2.00 g sample of gas has a pressure of  $3.00 \times 10^3$  mm Hg, a volume of  $1.95 \times 10^4$  mL, and a temperature of 455 K. What will be its pressure in atmospheres if the temperature is changed to 325 °C and volume is changed to 7.00 L?

13. How many moles and how many molecules of a gas are in a 255 ml aerosol can at a pressure of 855 torr and a temperature of 48 °C?

14. Some commercial drain cleaners contain two components: sodium hydroxide and aluminum powder. When the mixture is poured down a clogged drain, the following reaction occurs:



The heat generated in this reaction helps to melt away obstructions such as grease.

a. Calculate the volume of H<sub>2</sub> formed at STP by the reaction of 3.12 g of Al with NaOH?

15. What is the molar mass of a gas if a 1.50 L sample of the gas at 25 °C and 745 mm Hg has a mass of 2.89 g

16. Calculate the density for CH<sub>4</sub>, (methane) at 0 °C and 1.00 atm.



17. Classify the intermolecular forces between molecules of each of the following liquids.

- a. CO
- b. O<sub>2</sub>
- c. CH<sub>3</sub>OH
- d. HF
- e. CO<sub>2</sub>
- f. CH<sub>2</sub>O

18. Circle the correct answer for the following

- a. The higher boiling point: N<sub>2</sub> or CO?
- b. Weaker intermolecular forces: H<sub>2</sub>O or H<sub>2</sub>S?
- c. Lower vapor pressure: NH<sub>3</sub> or IF?
- d. Lower boiling point SO<sub>2</sub> or HF

19. Calculate the molar mass of 8.00 g of gas with a volume of 12.0 L at STP

20. If 40.0 g of acetylene[C<sub>2</sub>H<sub>2</sub>] undergoes combustion...

The balanced equation is:

a. How many liters of oxygen are required to burn the acetylene if the reaction takes place at 155 °C and 2.60 atm

b. How many mLs of carbon dioxide are produced at STP?

21. Show the polarity of the following molecules:

a. HBr

b.  $\begin{array}{c} \text{H}-\text{S} \quad | \\ | \\ \text{H} \end{array}$

c. S=C=S

22. a. Give the properties of gases, liquids and solids.

b. Explain the differences in the strength of the attractive forces between molecules in the gas, liquid and solid phase.

